

# James Aitken

MB, BS; LRCP, MRCS; MS; FRCS (Edin); FCS(SA); FRACS

## General and Colorectal Surgery

Unit 4,  
77 Grand Boulevard,  
Joondalup, 6027

Suite 41  
Hollywood Medical Centre,  
85 Monash Avenue,  
Nedlands, 6009

Tel: 6389 0244

Fax: 6389 0255

[www.perthcolorectal.com.au](http://www.perthcolorectal.com.au)

email: [info@perthcolorectal.com.au](mailto:info@perthcolorectal.com.au)

**All correspondence to Hollywood**

## RESECTION FOR RIGHT SIDED COLON CANCER.

This general guide is designed to provide background information to the operation that you will shortly undergo. It aims to supplement verbal discussion, to answer common questions and to be readily available as an *aide memoir*. It cannot cover in detail every aspect of your individual operation and may not deal with some areas that are of particular concern to you. These can be dealt with individually.

You should feel free to ask about any aspect of your care. All your questions will be answered fully, honestly and in as much detail as you wish. In the heat of the moment it is easy for questions that you intended to ask to slip from your mind. You should note on paper any questions that you may have.

Further information is available at the web site above. This site also provides links to other sites that may provide additional information.

### What is involved?

The aim of the operation is to remove the colon and small bowel that contains the Crohn's disease. At the same time the blood vessels and tissues supporting the bowel are also removed. A general examination will then be performed to determine the exact extent of the Crohn's disease and to ensure there are no other abnormalities in the abdomen. The affected colon will then be resected and the two ends joined together. The wound is then closed. There may or may not be skin sutures that have to be removed some 8-10 day later

### Open or laparoscopic (keyhole) surgery?

Large bowel surgery has traditionally been by open surgery. The incision was either a low transverse incision or an 'up and down' incision in the middle of the abdomen. It is now possible to offer selected patients laparoscopic (key hole) surgery. Laparoscopic surgery means the incisions are small, but once inside the abdomen a laparoscopic operation is almost identical to open surgery. That is why it is often called 'minimal access surgery'. The advantage of laparoscopic surgery is that the post-operative recovery is quicker.

Patients having this type of surgery will normally be offered laparoscopic surgery. One incision will have to be larger than the others in order to extract the bowel. Some operations that commence laparoscopically cannot be completed and have to be converted to an open operation. The operation is then completed in the normal manner.

For the avoidance of confusion the remainder of this document assumes your operation will be open surgery. If your operation is undertaken laparoscopically your management will be the same, but progress will be faster and discharge earlier.

### Before the operation.

The anaesthetist will assess you and will discuss the various ways of controlling post-operative pain (see below). Prior to the operation you may be given a bowel preparation to empty and clean the bowel.

It is important that we know every medical issue that might affect you. What may appear unimportant to you may be essential for us to know. In particular, we need to know all the drugs you are taking. You should bring them to the hospital in their original packet. Unless advised specifically to the contrary you should take all your drugs up to and including the morning of surgery. The exceptions to this are blood thinning agents, such as aspirin, plavix or warfarin, and diabetic drugs. These require special arrangements and must be discussed on an individual basis.

Once you are asleep various tubes and lines will be inserted into your veins, the bladder and through your nose into your stomach. These will be removed during the first 2-4 days following surgery.

### **Will a colostomy be required?**

A stoma (colostomy or ileostomy) is when part of the bowel is brought up onto the abdominal wall and a bag has to be worn. A stoma is not required as part of this operation.

### **Pain relief.**

Proper pain relief is very important for both holistic and physiological reasons. Your post-operative recovery will be slower if you do not have adequate pain relief. Patients often have an understandable reluctance to take pain relieving drugs. This is a mistake and may increase post-operative complications. The principal that underlies all methods of pain relief is that the drugs work best if you anticipate the pain. A small quantity of the drug taken regularly (even if pain free at that time) will work better than waiting for the pain to occur and then taking a larger dose.

Before the operation the anaesthetist will offering you two types of pain relief. The first is an epidural anaesthetic. This requires a needle to be inserted into your back and drugs are given through a fine catheter. The alternative is 'Patient Control Analgesia (PCA). With this technique you press a button as and when you feel the pain and a small dose of the pain relieving drug is administered. The advantage of these techniques is that a small quantity of the pain relieving drug can be administered on an on-going or regular basis and prevent the pain rather than treat it after it occurs. This is by far the most effective form of pain relief. Alternatively, regular injections can be administered. This is not as effective as they are usually administered after the pain has occurred.

After a couple of days adequate pain relief can normally be achieved by oral medication. Regular panadol, regardless of whether you have pain or not, is the foundation drug. If panadol alone is not adequate, additional, stronger painkillers and/or anti-inflammatory drugs can be taken on top of the panadol for break through pain. You should use panadol for background pain relief for a week after your operation. Many strong painkilling medications contain morphine, codeine or a derivative of these drugs. One of the side effects of these drugs is constipation.

### **The second 24 hours.**

You may be attached to various monitors and numerous observations will be performed. The physiotherapist will visit you and will ensure your lungs are clear and free of secretions and exercise your legs. You will be sat in a chair during part of the day.

### **Days three to five.**

The tubes may be removed. You will be permitted fluid by mouth and you may pass some flatus (wind) through the anus.

### **Days five to ten.**

You will start eating, moving around the ward, have a bath and generally return to normal, but limited, activities. Your bowel will start to work but maybe a little erratic and you may have some diarrhoea. Patients recover at different speeds and you should not be concerned if your progress appears slower than you anticipated.

### **Going home.**

You will normally return home 7 - 10 days after your surgery, but this will vary with your progress and home circumstances. If your right hemicolectomy is completed laparoscopically you will normally be discharged in 4 – 7 days. You will obviously be tired and you should plan to rest during each day. You should avoid domestic activities for at least the first three weeks. Sitting in a high backed chair can reduce the strain on your abdominal wound as it is easier to get up out of.

### **Recovery over the first six months.**

Over the first six weeks you should gradually increase the exercise you take. At six weeks you will be about 80% recovered (95% if laparoscopic). To fully recover to the point that you feel you have not had an operation will take three to six months, depending on your age and pre-operative fitness.

You should avoid strenuous exercise for four weeks. 'Little but often' should be your aim and a short walk two or three times a day is better than one long walk. Gradually increase the distance you walk over the next few weeks. The surgical wound has almost returned to full strength at ten days so it is almost impossible to 'over exercise' yourself to the extent that you damage the surgical area. If you feel comfortable doing a particular activity then it is very unlikely you will do yourself any harm. In general it is sudden, unplanned movements that cause problems.

As you start to feel better the likelihood is that you will overdo it and at this stage a couple of days of feeling well (and overdoing it) will be followed by a bad day as your body compensates. You have been warned! In general you should stop if you feel tired or if you feel pain.

For medico-legal reasons you must not drive for four weeks.

### **Bathing and showering.**

It is quite safe to get your wound wet with a shower or quick bath two or three days after your operation. However, long soaking baths with a Jeffery Archer novel should be avoided for at least three weeks as the wound will become soft and the scab may become infected. Adding salt to the bath will not help heal the wound and may make your skin dry and tight. After washing, pat the wound dry with a clean towel. A bath mat helps prevent slipping and a towel hooked around the bath taps can be a helpful lever when you try to get out. It can also be reassuring to have someone else in the house the first time you have a bath, even if you do not need help.

### **Sleep.**

Changes in your routine, restricted movement, lack of exercise and wound discomfort will interrupt your normal sleep pattern or wake you during the night. Uninterrupted sleep is more valuable than 'cat-napping' so you may find it helpful to take a pain killer before you go to bed.

You can resume sexual activity when this feels comfortable.

### **Eating.**

Your appetite will not be good for some weeks after surgery and you may feel aches, bloating and indigestion after meals. These symptoms usually disappear as you become

more active. You should take small, frequent meals with a good intake of protein (lean meat, dairy produce, fish *etc.*). A small amount of alcohol can improve your appetite and is not usually harmful.

### **Work.**

Your return to work depends on many factors, including your occupation, age and general health. You will definitely require one month off work, and some may require two months. It is better to feel completely well before you return to work rather than have to take more time off a few weeks or days later because you have returned to work too early.

### **The wound.**

A major concern to patients is that they will strain the wound and that it will rupture. With today's suture materials this is very unlikely. On the very few occasions that a wound does rupture it will be before you leave hospital. This would require an operation to repair the rupture. Once you have gone home a rupture is almost unheard of. If you 'over do it' the worst that will happen is that the wound will be very sore.

Wounds progress through several stages of healing. You may experience:-

- unusual tingling, numbness or itching sensations.
- a slightly hard or 'lumpy' feeling as new tissues form.
- pulling around the stitches or staples as the wound heals.

This is normal. Do not pull at any scabs as they act as a natural dressing and protect the new skin underneath. They will fall off when no longer required. You should seek help if any of the following occur:-

- the wound pain increases
- the wound becomes more reddened or swollen
- there is any discharge from the wound

### **Your post-operative bowel habit.**

This operation removes a segment of right colon. You may notice your stools are more frequent and a little loose immediately after the operation. It is likely to be diet related. This improves over the first three months and even further over the next six months. Some patients notice an ongoing improvement for up to two years. Depending on your individual circumstances it may be necessary to prescribe some tablets to help you.

Detailed information on how to improve your post-operative bowel habit is available on a separate advice sheet.

### **Surgical trainees**

Some patients may have their right hemicolectomy undertaken by a surgical trainee. A trainee performing a right hemicolectomy is normally, but not always, under the direct supervision of the consultant. It is important that, as part of their training, trainees gain independent experience whilst consultant cover is still immediately available. There is a substantial body of surgical literature that shows the outcome of operations undertaken by properly supervised trainees is no worse than those performed by the consultant. This literature specifically includes right hemicolectomy.

### **Follow-up.**

You will be reviewed about six weeks after your surgery. Further follow up will depend on many factors and is individualised for each patient. Further colonoscopies may be performed.

If you have any concerns between routine follow-up visits do not delay until your next scheduled appointment. Make an earlier appointment.

## What can go wrong?

You will be undergoing a major operation. Major operations are sometimes complicated by adverse events. That said, the surgeons, anaesthetists and nurses caring for you have an extensive experience in what is, to them, a routinely performed operation.

However, it is important that you appreciate that major surgery does carry risk and complications can and do occur. The table below summarises the potential risks and complications. This table is not exhaustive and if you have any concerns you should ask before you sign the consent form. It is not intended to alarm as most patients will not have a serious complication. However, one third of patients having a colonic resection will have an event that delays their recovery. Although everything possible will be done to prevent the development of any complication, it is only possible to reduce, not eliminate, these events.

Patients with Crohn's Disease have a complication rate that is greater than the same operation undertaken for other conditions. This is because Crohn's Disease is a chronic condition that may affect patients in subtle ways – for example, their albumin may be low. Also patients with Crohn's Disease usually have their operation when medical therapy has failed, so they are less well, are often on drugs, such as steroids, that slow healing.

The majority of complications tend to occur in two groups. The first group are a direct effect of that particular operation. The most important specific complication of this operation is a leak where the bowel was joined (the anastomosis). This is a serious complication (2% chance) and usually requires a second operation as an emergency. This second operation might require an ileostomy. It may be possible to close this ileostomy at a second operation some months later.

The second group are general complications that can occur after any operation. The risk of these complications is greatly influenced by any pre-existing medical conditions. Examples include a previous heart attack, a chronic illness such as diabetes, and smoking.

The best way to manage potential complications is to prevent them occurring in the first place. Hence the use of preventative, or prophylactic, treatment. It is important to correct any underlying medical conditions. For this reason it is essential you advise your doctors of all earlier operations and previous or ongoing medical illness. All your drugs should be brought to the hospital and shown to your doctors.

Risk	What happens	What may be done (options)
<i>General complications that may occur after any surgery</i>		
Clot in legs (DVT)	A clot forms in the legs. This may make the legs swell. The clot may break away into the lungs. This is a pulmonary embolus.	Blood thinning drugs (heparin) started at the time of surgery. TED stockings.
Post-operative bleeding	Blood leaks into the abdomen or out through a drain	1. blood transfusion 2. re-operation
Wound infection	An infection, including the development of pus, occurs in the wound	Antibiotics started at the time of surgery. Drainage of any pus is required, and this may require another operation or drainage under radio-logical guidance
Chest infection	A pneumonia develops	Antibiotics are required. A few patients require ventilation (in ICU)
Wound dehiscence	The wound opens up	Surgical repair within a few hours.
Incisional hernias through the wound	A weakness develops in the wound. The bowel then slips through the abdominal wall and a bulge appears. This usually occurs more than six months after surgery. The overall life	A surgical repair, usually with mesh, is required.

time risk is >15%. This risk is increased in patients who have either chemo-therapy or radiotherapy, a wound infection or a chronic illness like diabetes or obesity.

Urinary tract infection	Bacteria enter the bladder	Antibiotics
Bladder may not empty	It is not possible to pass urine. As the bladder get full, the patient gets uncomfortable.	The catheter is re-inserted and removed a few days later. Normally this solves the problem. Sometimes a catheter is required for 2-3 weeks. In men, prostate surgery may be required.
Vascular event	Stroke Heart attack	Each event managed on its own merits. Normally a period in ICU is required.
Death		In a fit person <2%. In other patients the risk is very dependent on pre-existing medical problems. A general estimate can be made by the use of risk prediction tables.

#### *Complications that may occur after bowel surgery*

Anastomotic leak	The join between the two ends of the bowel develops a leak. In this operation <2%.	<ol style="list-style-type: none"> <li>1. antibiotics alone</li> <li>2. Drainage under radiological guidance</li> <li>3. further surgery, including an stoma if not already present</li> </ol>
Post operative ileus	The bowel remains paralysed for longer then the usual 3-4 days	<ol style="list-style-type: none"> <li>1. a tube through the nose is inserted/left in the stomach</li> <li>2. various drugs may be given</li> <li>3. although it normally resolves in 3-5 days an ileus can occasionally be so prolonged that intra-venous feeding (TPN) is required.</li> </ol>
Bowel blockage (adhesions)	Scar tissue in the abdomen blocks the bowel. This can occur within a few days of surgery, or many years latter (or any time in between). The life time risk is 15-20%	A NGT and IVI settles most. Some patients require further surgery.

<b><i>What increases the risk of surgery</i></b>	<b><i>Examples</i></b>	<b><i>Why is the risk increased</i></b>
Medical illness	Pre-existing general medical conditions such as endocrine disorders, heart attacks or strokes etc.	As far as possible pre-existing medical problems will be corrected prior to surgery
Previous surgery		Scarred tissue is normally of poor quality and does not heal well
Obesity	All risks increased	<ol style="list-style-type: none"> <li>1. poor quality tissue</li> <li>2. poor mobilisation leading to increased risk of DVT, chest infection</li> <li>3. poor blood supply so the risk of wound or anastomotic failure is much increased</li> <li>4. extra strain on the wound, heart etc</li> </ol>
Drugs	Examples include steroids, aspirin, blood thinning agents	Normally because they increase the risk of bleeding, infection or decrease the quality of wound healing
Diabetes		<ol style="list-style-type: none"> <li>1. ability to combat infection reduced</li> <li>2. poor blood supply</li> <li>3. slow healing</li> </ol>
Smoking		Increased risk of anastomotic leak, infection,, vascular events and thrombosis

---

**Definitions**

IVI	Intravenous infusion ('a drip')	
NGT	Nasogastric tube	A fine tube from through the nose into the stomach to drain the stomach and stop vomiting.
ICU	Intensive Care Unit	For very ill patients, or those requiring ventilation
Ventilation		Placing patients on a machine that does the breathing for them. A tube is placed through the mouth into the upper airway.

---